

Physiotherapy and occupational therapy and mild to moderate parkinson disease-reply

Clarke, Carl E; Walker, Marion F; Sackley, Catherine M

DOI:

[10.1001/jamaneurol.2016.1283](https://doi.org/10.1001/jamaneurol.2016.1283)

License:

None: All rights reserved

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Clarke, CE, Walker, MF & Sackley, CM 2016, 'Physiotherapy and occupational therapy and mild to moderate parkinson disease-reply', *JAMA Neurology*. <https://doi.org/10.1001/jamaneurol.2016.1283>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked 14/06/2016

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Letters

COMMENT & RESPONSE

In Reply On behalf of the PD REHAB Collaborative Group, we are pleased that the PD REHAB Trial¹ has prompted debate. The issues raised are common to the 3 letters²⁻⁴ and all were covered in the original article.¹

Eligibility was based on the uncertainty principle. Investigators were mostly uncertain of the value of physiotherapy and occupational therapy in patients with mild to moderate Parkinson disease (PD). So, the results of the trial can only be applied to patients with mild to moderate disease. This is not in conflict with our Cochrane review of physical therapies in PD, which showed short-term benefits in motor function and activities of daily living but, importantly, not in quality of life in more severe disease.⁵

The Nottingham Extended Activities of Daily Living Scale was used as the primary outcome so that any benefits could be compared with those in other disease areas where it has been used before (ie, stroke and aging). It has been used in previous PD trials and correlated well with the Unified Parkinson's Disease Rating Scale activities of daily living scale in our pilot PD/OT trial.⁶ This approach precluded the use of an individualized outcome as suggested by de Vries et al⁴ in their letter. Although the Nottingham Extended Activities of Daily Living Scale does not directly address gait and transfers, it does assess issues such as walking outside, using public transport, and climbing stairs, which are of more practical use and more important to patients. Indeed, our patient advisory group supported the use of the Nottingham Extended Activities of Daily Living as the primary outcome for this very reason.

We too were surprised by the low dose of both therapies, but this is what is being delivered to patients with mild to moderate PD in the National Health Service (NHS) today. We do not agree with Mestriner³ that this is nonadherence; it is a low-dose intervention. We were informed by therapists before the trial that both physiotherapists and occupational therapists see such referrals, so they wanted the combined design of the PD REHAB Trial. A large number of centers were deliberately secured to aid recruitment and to ensure that the results could be generalized across the NHS. Therefore, PD REHAB is a pragmatic trial of standard care in the NHS and not a study of a novel high-intensity form of therapy. This is the design that the funder, the NHS Health Technology Assessment Programme, wanted to inform deci-

sions to be made about the future delivery of such services within the NHS.

The PD REHAB Trial is a high-quality, precise, adequately powered study that demonstrates that low-dose, patient-centered, goal-directed physiotherapy and occupational therapy in early PD is ineffective. We cannot ignore or seek to discredit important results that we do not find palatable. We owe it to our patients to deliver the best care possible, so future research should explore the development and testing of more structured and intensive physical and occupational therapy programs in patients with all stages of PD.

Carl E. Clarke, MD

Marion F. Walker, PhD

Catherine M. Sackley, PhD

Author Affiliations: School of Clinical and Experimental Medicine, College of Medical and Dental Sciences, University of Birmingham, Birmingham, England (Clarke); Department of Neurology, Sandwell and West Birmingham Hospitals NHS Trust, City Hospital, Birmingham, England (Clarke); University of Nottingham, Nottingham, England (Walker); King's College London, London, England (Sackley).

Corresponding Author: Carl E. Clarke, MD, Department of Neurology, Sandwell and West Birmingham Hospitals NHS Trust, City Hospital, Dudley Road, Birmingham B18 7QH, England (carlclarke@nhs.net).

Published Online: May 31, 2016. doi:[10.1001/jamaneurol.2016.1283](https://doi.org/10.1001/jamaneurol.2016.1283).

Conflict of Interest Disclosures: None reported.

Additional Contributions: We are grateful to Natalie Ives, BSc, MSc (senior statistician, University of Birmingham Clinical Trials Unit), for her critical review and comments on this reply. She did not receive compensation.

1. Clarke CE, Patel S, Ives N, et al; PD REHAB Collaborative Group. Physiotherapy and occupational therapy vs no therapy in mild to moderate Parkinson disease: a randomized clinical trial. *JAMA Neurol*. 2016;73(3):291-299.
2. Ellis T, Tickle-Degnen L, Saint-Hilaire M. Physiotherapy and occupational therapy and mild to moderate parkinson disease [published online May 31, 2016]. *JAMA Neurol*. doi:[10.1001/jamaneurol.2016.1274](https://doi.org/10.1001/jamaneurol.2016.1274).
3. Mestriner RG. Physiotherapy and occupational therapy and mild to moderate parkinson disease [published online May 31, 2016]. *JAMA Neurol*. doi:[10.1001/jamaneurol.2016.1280](https://doi.org/10.1001/jamaneurol.2016.1280).
4. de Vries NM, Sturkenboom IH, Bloem BR. Physiotherapy and occupational therapy and mild to moderate parkinson disease [published online May 31, 2016]. *JAMA Neurol*. doi:[10.1001/jamaneurol.2016.1277](https://doi.org/10.1001/jamaneurol.2016.1277).
5. Tomlinson CL, Patel S, Meek C, et al. Physiotherapy versus placebo or no intervention in Parkinson's disease. *Cochrane Database Syst Rev*. 2013;9:CD002817.
6. Clarke CE, Furlston A, Morgan E, et al. Pilot randomised controlled trial of occupational therapy to optimise independence in Parkinson's disease: the PD OT Trial. *J Neurol Neurosurg Psychiatry*. 2009;80(9):976-978.